



Endpoints

**in Alzheimer's disease
clinical trials**

An educational overview
of clinical trial endpoints
currently used in
Alzheimer's disease

Co-created with the clinical and research community in mind

Prepared June 2022

M-XX-00009769

Introduction

Alzheimer’s disease (AD) is a progressive neurodegenerative disease characterized by cognitive, functional, and behavioral deficits. Our understanding of the causes and potential treatments for AD is continually evolving. The goal of this interactive brochure is to provide greater recognition of what clinical trials measure (endpoints) in AD. We also aim to describe the relevance of common AD endpoints from the perspective of clinical teams managing people with AD.

Those who may find this guide useful include neurologists, psychiatrists, primary care physicians, early-career clinicians, and other health care professionals who want to understand the endpoints used in AD clinical trials. It is also suitable for trial investigators and staff involved in the design, management, and administration of clinical trials in AD. Information about regulatory guidance, statistical considerations, and biomarkers has been included in an interactive and approachable way.

We have supplemented this information with our years of research and clinical experience, adding our commentary and insights about the value and utility of the methods that investigators use to measure and understand AD research. *Where you see a speech bubble, this indicates where we have added our perspective.*



Richard S. Isaacson, M.D., Director of the Alzheimer’s Prevention Clinic at Weill Cornell Medicine and New York-Presbyterian, New York, USA.



Ramin Nilforooshan, M.D., MRCPsych., Consultant Psychiatrist, Surrey and Borders Partnership NHS Foundation Trust, Chertsey, UK and Visiting Professor at the University of Surrey, UK. Clinical Lead for the UK Dementia Research Institute CR&T.



Kathleen Welsh-Bohmer, Ph.D., Professor of Psychiatry and Behavioral Sciences; Professor in Neurology and Faculty Network Member of the Duke Institute for Brain Sciences, Duke University School of Medicine, Durham, North Carolina, USA.

This booklet is for educational purposes only and is not intended to be a comprehensive assessment of clinical trial endpoints in Alzheimer’s disease. The information included here along with commentary and input from members of the scientific community does not represent an endorsement or recommendation of any endpoints.

Contents

Endpoints in AD clinical trials

Individual clinical outcome assessments and endpoints summarized

Clinical Dementia Rating-Sum of Boxes (CDR-SB)
Alzheimer's Disease Assessment Scale-Cognitive Subscale (ADAS-Cog)
Alzheimer's Disease Cooperative Study-Activities of Daily Living (ADCS-ADL)
Mini-Mental State Examination (MMSE)
Verbal Fluency Task (VFT) Score
Wechsler Adult Intelligence Scale, Fourth Edition (WAIS-IV) – Coding Test
Repeatable Battery for the Assessment of Neuropsychological Status (RBANS)
Preclinical Alzheimer Cognitive Composite 5 (PACC5)
Disability Assessment for Dementia (DAD)
Functional Activities Questionnaire (FAQ)
Amsterdam - Instrumental Activities of Daily Living (A-IADL)
Quality of Life in Alzheimer's Disease (QoL-AD)
5-Level EuroQoL-5D (EQ-5D-5L)
Alzheimer's Disease Composite Score (ADCOMS)
Cognitive Function Instrument (CFI)
Integrated Alzheimer's Disease Rating Scale (iADRS)
Alzheimer's Disease Cooperative Study-Clinical Global Impression of Change (ADCS-CGIC)
Neuropsychiatric Inventory-Questionnaire (NPI-Q)
Columbia-Suicide Severity Rating Scale (C-SSRS)
Zarit Burden Interview (ZBI)
Resource Utilization in Dementia-Lite Version (RUD-Lite)

Fluid-based and imaging biomarkers in AD clinical trials

Individual biomarkers summarized


CSF – amyloid beta ($A\beta$), CSF – tau, CSF – amyloid beta and tau ratios,
Amyloid PET, Tau PET, Volumetric MRI

References



Index of clinical outcome assessments (COAs) and endpoints included in this booklet,* by category

Several types of COAs and endpoints are used in AD



*This is not a comprehensive list of AD endpoints

Also considered as a **Proxy-reported outcome, which is an assessment in which someone other than the patient reports on patient symptoms.¹

ADAS-Cog, Alzheimer's Disease Assessment Scale-Cognitive Subscale; ADCOMS, Alzheimer's Disease Composite Score; ADCS-ADL, Alzheimer's Disease Cooperative Study - Activities of Daily Living; ADCS-CGIC, Alzheimer's Disease Cooperative Study-Clinical Global Impression of Change; A-IADL, Amsterdam - Instrumental Activities of Daily Living; CDR-SB, Clinical Dementia Rating-Sum of Boxes; CFI, Cognitive Function Instrument; CSF, cerebrospinal fluid; C-SSRS, Columbia-Suicide Severity Rating Scale; DAD, Disability Assessment for Dementia; EQ-5D-5L, 5-Level EuroQoL-5D; FAQ, Functional Activities Questionnaire; iADRS, Integrated Alzheimer's Disease Rating Scale; MMSE, Mini-Mental State Examination; MRI, magnetic resonance imaging; NPI-Q, Neuropsychiatric Inventory-Questionnaire; PACC5, Preclinical Alzheimer Cognitive Composite 5; PET, positron emission tomography; QoL-AD, Quality of Life in Alzheimer's Disease; RBANS, Repeatable Battery for the Assessment of Neuropsychological Status; RUD-Lite, Resource Utilization in Dementia – Lite Version; WAIS-IV, Wechsler Adult Intelligence Scale – Fourth Edition; ZBI, Zarit Burden Interview.


Endpoints in AD clinical trials

Introduction

Endpoints are precisely defined variables that are intended to reflect an outcome of interest that will address a specific research question.² In clinical trials, endpoints are used to evaluate treatment efficacy and/or safety.


COAs differ from endpoints, in that they are the instruments used to evaluate the intended outcome. They can be made through reports by patients, clinicians, or non-clinician observers.² There are four main types of COA (Figure 1).²

Figure 1: Types of COA²



Clinical trials employ a hierarchy of predefined endpoints (Figure 2).³ In AD clinical trials, it is common to look at multiple endpoints. However, in statistical terms, a greater number of endpoints contribute to a potential reduction in study power, i.e. the likelihood or ability of a study to detect a true treatment effect.³ Therefore, determining which endpoints to include in an AD trial requires a balance between trying to best capture the disease manifestations and ensuring the study is adequately powered.

Figure 2: The hierarchy of predefined endpoints in clinical trials³




AD is a multifactorial disease with an evolving understanding of pathogenesis and treatment

AD is characterized by a progressive decline in cognitive function and impaired activities of daily living, leading to diminished quality of life.^{4,5} Endpoints in AD are specific COAs of cognition, function, overall clinical response (i.e. global assessment), behavior and psychiatric symptoms, or quality of life (**Figure 3**).⁶ Endpoints contribute to the interpretation of clinical efficacy of treatments in AD,⁷ and they must be sensitive to change (to detect treatment effects) and clinically meaningful.⁸

People with dementia and family caregivers have stated that:⁸

- Questionnaire content for endpoints should be clear and limited in volume
- Cognition should be a core endpoint but should be used in conjunction with others to provide contextual, qualitative information

Figure 3: Domains of AD outcome assessment⁶




AD, Alzheimer's Disease

Regulatory guidance on primary endpoints in AD clinical trials

Regulatory agencies have provided guidance on the use of AD endpoints by disease stage (**Figure 4**).^{6,9} The US Food and Drug Administration (FDA) guidance states that a composite (combined cognitive and functional) outcome may serve as a primary endpoint in prodromal AD.¹⁰ Both the FDA and European Medicines Agency (EMA) guidelines emphasize the importance of incorporating the patient voice in developing acceptable measures that are clinically meaningful.¹¹

Figure 4: Regulatory guidance on disease stage applicability of endpoints in AD^{6,9}



^aAssessed by sensitive neuropsychological measures; ^bHowever, cognitive testing will likely be included in future regulatory guidance, for use in the preclinical stage. AD, Alzheimer's Disease; MCI, mild cognitive impairment.

Table 1. Selected primary and secondary endpoints used in AD clinical trials (marked by X)


Note: This is not a complete list of all endpoints used in these trials, and biomarker/imaging endpoints are not included in this table.

	GRADUATE I & II ^{12,13}	LAURIET ¹⁴	TAURIEL ¹⁵	EMERGE & ENGAGE ^{16,17}	AHEAD 3-45 ¹⁸	Clarity AD ¹⁹	ALZ-801 ^{*,20}	TRAILBLAZER -ALZ2 ²¹
CDR-SB	X (primary)	X	X (primary)	X (primary)		X (primary)		X (primary)
ADCOMS						X		
ADAS-Cog	X	X (primary)	X	X		X	X (primary)	X
MMSE	X	X		X				X
Verbal Fluency Task	X							
WAIS-IV – Coding	X							
RBANS			X					
PACC5					X (primary)			
iADRS								X
ADCS-ADL	X	X (primary)	X	X				X
DAD								
FAQ	X							
A-IADL			X					
QoL-AD								
EQ-5D-5L								
CFI					X			
ADCS-CGIC								
NPI-Q								
C-SSRS	X							
ZBI								
RUD-Lite								

*Secondary endpoints for the planned ALZ-801 phase 3 trial are noted to be function, activities of daily living, and behavior²⁰

AD, Alzheimer’s Disease; ADAS-Cog, Alzheimer’s Disease Assessment Scale-Cognitive Subscale; ADCOMS, Alzheimer’s Disease Composite Score; ADCS-ADL, Alzheimer’s Disease Cooperative Study - Activities of Daily Living; ADCS-CGIC, Alzheimer’s Disease Cooperative Study-Clinical Global Impression of Change; A-IADL, Amsterdam - Instrumental Activities of Daily Living; CDR-SB, Clinical Dementia Rating-Sum of Boxes; CFI, Cognitive Function Instrument; C-SSRS, Columbia-Suicide Severity Rating Scale; DAD, Disability Assessment for Dementia; EQ-5D-5L, 5-Level EuroQoL-5D; FAQ, Functional Activities Questionnaire; iADRS, Integrated Alzheimer’s Disease Rating Scale; MMSE, Mini-Mental State Examination; NPI-Q, Neuropsychiatric Inventory-Questionnaire; PACC5, Preclinical Alzheimer Cognitive Composite 5; QoL-AD, Quality of Life in Alzheimer’s Disease; RBANS, Repeatable Battery for the Assessment of Neuropsychological Status; RUD-Lite, Resource Utilization in Dementia - Lite Version; WAIS-IV, Wechsler Adult Intelligence Scale – Fourth Edition; ZBI, Zarit Burden Interview.

Table 2. Selected COAs/endpoints, by applicability across the spectrum of disease severity*

PRECLINICAL AD	MCI/PRODROMAL AD, MILD AD, MODERATE AD	SEVERE AD
		
CSF amyloid beta & tau	CDR-SB	CDR-SB
PET amyloid beta & tau	ADAS-Cog	MMSE
Volumetric MRI	MMSE	ADCS-ADL
PACC5	Verbal Fluency	DAD
CFI	WAIS-IV – Coding	FAQ
	RBANS	A-IADL
	PACC5	QoL-AD
	ADCS-ADL	EQ-5D-5L
	DAD	ADCOMS
	FAQ	ADCS-CGIC
	A-IADL	NPI-Q
	QoL-AD	C-SSRS
	EQ-5D-5L	ZBI & RUD-Lite
	ADCOMS	
	iADRS	
	ADCS-CGIC	
	NPI-Q	
	C-SSRS	
	ZBI & RUD-Lite	

*Categorization includes conclusions based on type of endpoint (i.e. domain).

AD, Alzheimer's Disease; ADAS-Cog, Alzheimer's Disease Assessment Scale-Cognitive Subscale; ADCOMS, Alzheimer's Disease Composite Score; ADCS-ADL, Alzheimer's Disease Cooperative Study - Activities of Daily Living; A-IADL, ADCS-CGIC, Alzheimer's Disease Cooperative Study-Clinical Global Impression of Change; Amsterdam - Instrumental Activities of Daily Living; CDR-SB, Clinical Dementia Rating-Sum of Boxes; CFI, Cognitive Function Instrument; COA, clinical outcome assessment; CSF, cerebrospinal fluid; C-SSRS, Columbia-Suicide Severity Rating Scale; DAD, Disability Assessment for Dementia; EQ-5D-5L, 5-Level EuroQoL-5D; FAQ, Functional Activities Questionnaire; iADRS, Integrated Alzheimer's Disease Rating Scale; MCI, mild cognitive impairment; MMSE, Mini-Mental State Examination; MRI, magnetic resonance imaging; NPI-Q, Neuropsychiatric Inventory-Questionnaire; PACC5, Preclinical Alzheimer Cognitive Composite 5; PET, positron emission tomography; QoL-AD, Quality of Life in Alzheimer's Disease; RBANS, Repeatable Battery for the Assessment of Neuropsychological Status; RUD-Lite, Resource Utilization in Dementia - Lite Version; WAIS-IV, Wechsler Adult Intelligence Scale – Fourth Edition; ZBI, Zarit Burden Interview.

Individual endpoints summarized

In this section, we summarize several COAs/endpoints used in AD clinical trials, highlighting their focus, use in clinical practice and research, components, and scoring.

Table 3. COAs/endpoints that encompass both cognitive and functional domains

COA/endpoint	What is assessed	What is not assessed/limitations
Clinical Dementia Rating-Sum of Boxes (CDR-SB)	<ul style="list-style-type: none"> Memory, judgement, problem solving, home and hobbies, orientation, and personal care²² 	<ul style="list-style-type: none"> Dependent on the accuracy and consistent availability of the patient's informant; time-intensive²³
Alzheimer's Disease Composite Score (ADCOMS)	<ul style="list-style-type: none"> Composite of ADAS-Cog, CDR-SB, and MMSE²⁴ 	<ul style="list-style-type: none"> Not used in clinical practice because of complexity – measures must be statistically combined; time-intensive
Cognitive Function Instrument (CFI)	<ul style="list-style-type: none"> Questionnaire (i.e. subjective assessment) of memory decline, cognitive difficulties, and functional decline^{25–27} 	<ul style="list-style-type: none"> Lacks the sensitivity required for clinical trials
Integrated Alzheimer's Disease Rating Scale (iADRS)	<ul style="list-style-type: none"> Combines cognition and daily function scores from the ADAS-Cog and the ADCS-IADL^{28,29} 	<ul style="list-style-type: none"> Recently developed, thus limited evidence/evaluation of the endpoint to date
Alzheimer's Disease Cooperative Study-Clinical Global Impression of Change (ADCS-CGIC)	<ul style="list-style-type: none"> General performance, mental cognitive state, activities of daily living, and behavior^{30,31} 	<ul style="list-style-type: none"> Not designed for use in the clinical practice setting



AD, Alzheimer's Disease; ADAS-Cog, Alzheimer's Disease Assessment Scale-Cognitive Subscale; ADCS-IADL, Alzheimer's Disease Cooperative Study-Instrumental Activities of Daily Living; MCI, mild cognitive impairment; MMSE, Mini-Mental State Examination.

Table 4. COAs/endpoints evaluating cognition

COA/endpoint	What is assessed	What is not assessed/limitations
Alzheimer's Disease Assessment Scale-Cognitive Subscale (ADAS-Cog)	<ul style="list-style-type: none"> Memory, praxis, planning, and executive function^{32,33} 	<ul style="list-style-type: none"> Too lengthy for clinical practice,^{8,33} can be influenced by education/IQ, floor effects for MCI and early AD, may be insensitive to treatment effects
Mini-Mental State Examination (MMSE)	<ul style="list-style-type: none"> Severity of cognitive impairment in two sections: memory, orientation, and attention; and name, follow instructions, copy, and write.³⁴ Can be completed quickly and frequently 	<ul style="list-style-type: none"> No measure of executive function. Can be affected by sociocultural factors, age etc.³⁵ Most sensitive for moderate stages of dementia, less so for MCI^{5,35}
Verbal Fluency Task (VFT)	<ul style="list-style-type: none"> Semantic (e.g. name as many animals as you can) and phonemic fluency (e.g. name words beginning with F). Correlates with severity and risk of progression in AD³⁶ 	<ul style="list-style-type: none"> Premorbid IQ and socioeconomic factors impact results. Differences in language use between cultures limit applicability of findings to some populations
Wechsler Adult Intelligence Scale – Fourth Edition (WAIS-IV – Coding)	<ul style="list-style-type: none"> Executive function, processing speed, attention, short-term memory, and cognitive flexibility³⁷ 	<ul style="list-style-type: none"> Limited utility in patients with lower levels of cognition³⁸
Repeatable Battery for the Assessment of Neuropsychology Status (RBANS)	<ul style="list-style-type: none"> Short-term and delayed memory, visuospatial ability, language, and attention³⁹ 	<ul style="list-style-type: none"> Insensitive to later-stage clinical progress; time-intensive
Preclinical Alzheimer Cognitive Composite 5 (PACC5)	<ul style="list-style-type: none"> Composite of subscales from several measures designed for asymptomatic AD. Includes executive function and general cognition⁴⁰ 	<ul style="list-style-type: none"> Not used in clinical practice. As a combination of other measures, it may be time consuming

AD, Alzheimer's Disease; IQ, intelligence quotient; MCI, mild cognitive impairment.



Table 5. COAs/endpoints assessing function

COA/endpoint	What is assessed	What is not assessed/limitations
Disability Assessment for Dementia (DAD)	<ul style="list-style-type: none"> The ability to perform everyday activities, tasks, and behaviors. Basic and instrumental ADLs, and leisure activities are included⁴¹ 	<ul style="list-style-type: none"> Limited utility in early-stage AD
Functional Activities Questionnaire (FAQ)	<ul style="list-style-type: none"> Used to measure ADLs for the previous 4 weeks. A total score of ≥ 6 may be used to discern MCI from mild AD⁴² 	<ul style="list-style-type: none"> If patient has not performed an activity it cannot be assessed, leaving the questionnaire incomplete⁴²
Amsterdam - Instrumental Activities of Daily Living (A-IADL)	<ul style="list-style-type: none"> Aimed at detecting early-stage and early-onset dementia. Items cover household, personal finance, and leisure activities⁴³ 	<ul style="list-style-type: none"> The 70-item version is lengthy⁴³
Alzheimer's Disease Cooperative Study - Activities of Daily Living (ADCS-ADL)	<ul style="list-style-type: none"> Basic and instrumental ADL items (eating, walking, reading, cooking, etc.)⁴⁴ 	<ul style="list-style-type: none"> Functional impairment may not be detected in MCI,⁴⁵ relies on caregiver view and may be prone to bias,⁴⁴ time-intensive

AD, Alzheimer's Disease; ADL, activities of daily living; MCI, mild cognitive impairment.

Table 6. COA/endpoints measuring quality of life

COAs/endpoint	What is assessed	What is not assessed/limitations
Quality of Life in Alzheimer's Disease (QoL-AD)	<ul style="list-style-type: none"> QoL, physical condition, mood, interpersonal relationships, ability to participate in meaningful activity, and financial situation^{4,46} 	<ul style="list-style-type: none"> Caregiver-rated QoL-AD score is typically lower than self-rated QoL-AD.^{47,48} Self-assessed QoL-AD may have limited response and correlation to changes in clinical outcomes⁴⁷
5-Level EuroQoL-5D (EQ-5D-5L)	<ul style="list-style-type: none"> QoL of 5 domains: mobility, self-care, usual activities, pain/discomfort, and anxiety/depression⁴⁹ 	<ul style="list-style-type: none"> The self-completed EQ-5D-5L may not accurately reflect clinically important changes (e.g. functional assessment and CDR)⁵⁰

CDR, clinical dementia rating; QoL, quality of life.

Table 7. Psychiatric COAs/endpoints

COA/endpoint	What is assessed	What is not assessed/limitations
Neuropsychiatric Inventory–Questionnaire (NPI-Q)	<ul style="list-style-type: none"> Neuropsychiatric symptoms across 12 domains⁵¹ 	<ul style="list-style-type: none"> Does not include clinician rating
Columbia-Suicide Severity Rating Scale (C-SSRS)	<ul style="list-style-type: none"> Suicidal ideation and suicidal behavior: severity of ideation, intensity of ideation, behavior subscale, and lethality subscale⁵² 	<ul style="list-style-type: none"> Concerns regarding the appropriateness of use in patients with dementia; not specifically validated in the elderly population^{53,54}

**Table 8. COAs/endpoints on caregiving burden**

COA/endpoint	What is assessed	What is not assessed/limitations
Zarit Burden Interview (ZBI)	<ul style="list-style-type: none"> Caregiver burden (29 items including caregiver health, psychological well-being, relationship with the patient, social life, and finances)⁵⁵ 	<ul style="list-style-type: none"> Reflects caregiver burden but not directly related to severity of patient's disease⁵⁵
Resource Utilization in Dementia – Lite Version (RUD-Lite)	<ul style="list-style-type: none"> Cost-effectiveness, resource use for dementia care^{56,57} 	<ul style="list-style-type: none"> May not necessarily reflect resource use during clinical trials; resource use tends to be delayed



Clinical Dementia Rating–Sum of Boxes (CDR-SB)

The CDR-SB is a clinician-reported assessment of cognition and function via a semi-structured interview with the patient and informant (e.g. a caregiver or family member).^{22,58,59} It is suitable for assessment across the disease spectrum, from prodromal to moderate AD stages.²³ Limitations include length of assessment; dependency on accuracy and consistent availability of the patient's informant,²³ and risk of disease stage misclassification with moderate stage AD.⁵

Unlike the CDR-SB, the CDR-global scale (CDR-GS), is typically used for staging purposes, and characterizes a participant's level of impairment according to the following categories: 0 (normal), 0.5 (very mild/prodromal dementia), 1 (mild dementia), 2 (moderate dementia), and 3 (severe dementia).⁶⁰

Domains:²²

Memory, orientation, judgment and problem solving, community affairs, home and hobbies, and personal care

Each domain is assessed for degree of impairment on a 5-point scale:

0 = none; 0.5 = questionable; 1 = mild;
2 = moderate; 3 = severe

Sum of boxes:⁵⁸ sum of the ratings for each domain (maximum score of 18)

Proposed meaningful change threshold*
by disease stage:⁶¹ MCI: +1; mild AD: +2;
moderate-severe: +2

*A meaningful change threshold is the level of score change on an outcome that is perceived as meaningful in the target population. Proposed thresholds provided throughout this brochure are based on limited evidence from the literature, with individual studies adopting a variety of approaches/methodologies. Please refer to the specific references for details on the methods used and see the PFDD guidance 3 for recommended gold standard approaches⁶²

Insights on use in clinical practice and research settings**

- *Rarely used in a day-to-day clinical setting*
- *Relies heavily on caregivers and requires the same rater for every measurement*
- *Unlikely to provide a continuous measure for each domain; informants usually provide information that covers only 1–2 weeks of changes, not 3 months*

**This is the experience and thoughts of our expert authors



Alzheimer's Disease Assessment Scale–Cognitive Subscale (ADAS-Cog)

ADAS-Cog is a performance outcome and the most frequently used cognitive outcome in AD clinical trials. ADAS-Cog-11 assesses memory, praxis, concentration, planning, and executive function.^{32,33} Expanded versions include: ADAS-Cog-12: + delayed word recall; ADAS-Cog-13: + number cancellation; and ADAS-Cog-14: + maze completion.⁶³ Key limitations include duration of assessment (not routinely used in clinical practice)^{8,33} and its potentially limited ability to detect treatment effects as a single endpoint.⁶⁴

ADAS-Cog consists of the following subtests:⁶⁵

• Word recall	• Comprehension
• Following commands	• Word-finding ability
• Constructional praxis	• Spoken language ability
• Naming objects	• Remembering test instructions
• Ideational praxis	• Delayed word recall (ADAS-Cog-12)
• Orientation	• Number cancellation (ADAS-Cog-13)
• Word recognition	• Maze completion (ADAS-Cog-14)

The maximum score is 70 points; higher scores indicate more severe impairment

Proposed meaningful change threshold* in early AD: 3-point decline

*A meaningful change threshold is the level of score change on an outcome that is perceived as meaningful in the target population. Proposed thresholds provided throughout this brochure are based on limited evidence from the literature, with individual studies adopting a variety of approaches/methodologies. Please refer to the specific references for details on the methods used and see the PFDD guidance 3 for recommended gold standard approaches⁶²

Insights on use in clinical practice and research settings**

- *A common outcome for cognition in clinical trials but not used in clinical settings*
- *Lengthy, time-dependent, and requires the scale box*
- *Premorbid IQ/education may impact the assessment*
- *Score is not meaningful if it remains unchanged but there is a decline in daily function*
- *Best suited to mild/moderate disease – floor effects for MCI and early AD*

**This is the experience and thoughts of our expert authors



Alzheimer's Disease Cooperative Study - Activities of Daily Living (ADCS-ADL)

The ADCS-ADL is an observer-reported outcome that assesses basic and instrumental activities of daily living items.⁴⁴ The scale was designed specifically for use in AD clinical trials, and caregivers or other informants complete the questionnaire.⁴⁴ However, impairment in functioning may not be detected in patients with mild cognitive impairment (MCI).⁴⁵

ADCS-ADL consists of the following subtests:⁴⁴

• Eating	• Handles mail
• Walking	• Hobbies or games
• Toileting	• Watching television
• Bathing	• Uses telephone
• Grooming	• Conversation
• Dressing	• Turns off lights
• Chooses clothes	• Clears the table
• Travel outside home	• Uses household appliances
• Shopping	• Puts away belongings
• Keeping appointments	• Finds belongings
• Able to be left alone	• Disposes of litter
• Discuss current events	• Obtains a beverage
• Reading	• Preparing a meal/snack
• Writing	

Total score ranges from 0 to 78; higher scores represent better functional level; lower scores indicate greater impairment

Insights on use in clinical practice and research settings*

- *Not used in clinical setting but may have value as a 'one-off' measure*
- *Provides a meaningful measure of cognitive change and translational impact*
- *Time-intensive*
- *Relies on caregiver view and is prone to bias*
- *Some questions cover topics that may not have occurred between ratings*
- *Insensitive to changes at the beginning of the disease; not sensitive enough for short clinical trials*

*This is the experience and thoughts of our expert authors



Mini-Mental State Examination (MMSE)

The Folstein MMSE is a performance outcome that quantitatively assesses the severity of cognitive impairment.³⁴ The MMSE is commonly used for inclusion criteria in AD clinical trials. It is considered to be most accurate for classifying moderate stages of dementia.⁵ Limitations include potential impact by sociocultural variables, age, and education and it is less sensitive to subtle cognitive changes in patients with MCI.³⁵

The **MMSE** consists of 2 sections:³⁵

- **Section 1:** covers orientation, memory, and attention (21 points maximum)
- **Section 2:** tests ability to name, follow verbal and written commands, write a sentence spontaneously, and copy a complex polygon (9 points)

Lower scores indicate increasing cognitive impairment; a score of 23 or 24 is indicative of cognitive impairment or dementia^{66,67}

Thresholds for a clinically meaningful decline increase with AD severity⁶¹

Insights on use in clinical practice and research settings*

- Used by most clinical trials and commonly used in clinical practice
- Premorbid IQ and age affect score, and cut-off points vary
- Quick to use and can be completed more frequently than other scales
- SMMSE is better standardized and has been used in various settings
- Not related to the outcome of patients in the clinical setting
- No assessment of executive function
- Use has decreased due to licensing/copyright requirements

*This is the experience and thoughts of our expert authors



Verbal Fluency Task Score

Semantic and phonemic fluency tests are performance outcomes that may be used to assess cognitive function in AD.³⁶ Verbal fluency, particularly semantic fluency, has been shown to correlate with severity of AD.³⁶ Semantic memory, and therefore fluency, is impaired in patients with amnesic MCI.⁶⁸ In addition, verbal fluency is associated with risk of progression to clinical dementia.³⁶

Examples of verbal fluency tests:^{36,68}

- **Semantic (or category) fluency:** *name as many animals as possible in 60 seconds*
- **Phonemic fluency:** *name as many words as possible beginning with the letter F, in 60 seconds*

Insights on use in clinical practice and research settings*

- *Used widely in the clinical setting*
- *Only takes 2 minutes and no special training required*
- *Helpful in clinical trials and for patient outcomes*
- *Education and premorbid IQ impact the results*
- *The influence of socioeconomic factors and language must be considered when used clinically or in global/other cultural contexts*

*This is the experience and thoughts of our expert authors



Wechsler Adult Intelligence Scale – Fourth Edition (WAIS-IV) – Coding Subtest

The WAIS is a common and broadly used human intelligence test. The digit symbol substitution test (DSST, i.e. coding test) of the WAIS-R instrument assesses executive function (via processing speed).^{37,69} The WAIS coding test has been shown to be highly predictive of time to conversion to AD in patients with MCI.^{37,69} Its main limitation is that its utility is limited in patients with lower levels of cognition.³⁸

The **WAIS-IV – Coding subtest evaluates**.³⁷

- *Visuomotor processing speed*
- *Short-term visual memory*
- *Cognitive flexibility*
- *Ability to absorb new material*
- *Attention, concentration, and motivation*

Using a key that matches numbers to simple symbols, the patient copies symbols as quickly as possible, during a 2-minute period³⁷

A higher score represents better performance,³⁷ in a study of patients with MCI, the mean score was 40.8 (SD 12.7)⁶⁹

Insights on use in clinical practice and research settings*

- *Used for cases that require detailed memory assessment to differentiate MCI from AD*
- *Helpful for clinical trials*
- *Shows the patient's stage of disease*
- *Used in LD services*
- *Provides an SD, which is helpful*
- *It is not helpful as cognitive score drops to the medium level*

*This is the experience and thoughts of our expert authors



Repeatable Battery for the Assessment of Neuropsychology Status (RBANS)

The RBANS assesses cognition through a variety of tests.³⁹ Alternate versions of the test exist, to counteract practice effects (i.e. improved performance due to familiarity with the test). The measure was designed for use over a wide range of cognitive status, from normal to moderate AD.³⁹ The RBANS is for use in both clinical practice and research settings. Diagnostic accuracy for both clinical AD and MCI due to AD have been demonstrated.^{70,71}

The RBANS consists of 12 subtests, which yield 5 index scores:³⁹

- **Immediate memory:**

- *List learning*
- *Story memory*

- **Visuospatial constructional:**

- *Figure copy*
- *Line orientation*

- **Language:**

- *Picture naming*
- *Semantic fluency*

- **Attention:**

- *Digit span*
- *Coding*

- **Delayed memory:**

- *List recall*
- *List recognition*
- *Story recall*
- *Figure recall*

Scores are scaled within each index and for a total score; a higher score indicates better performance³⁹

Insights on use in clinical practice and research settings*

- *Used for small samples of patients referred to a neuropsychologist for detailed memory testing*
- *Useful in clinical trials where later-stage clinical progress is not pertinent*
- *Takes much longer than normal memory assessment*

*This is the experience and thoughts of our expert authors



Preclinical Alzheimer Cognitive Composite 5 (PACC5)

The Alzheimer's Disease Cooperative Study PACC is a composite performance outcome assessing cognition.⁴⁰ Components of 4 cognition endpoints are included, evaluating memory, executive function, and general cognition. The PACC was designed specifically for use as a primary endpoint for clinical trials in asymptomatic AD.⁴⁰ The PACC5 includes a fifth domain, category fluency (semantic memory).⁷²

The **PACC5** consists of:^{40,72}

- **Total Recall** score from the FCSRT
- **Delayed Recall score** on the Logical Memory Ila subtest of the WAIS
- **Digit Symbol Substitution Test** score on the WAIS
- **Total MMSE** score
- **Category fluency**

Scores are combined into a single composite total score⁴⁰

Insights on use in clinical practice and research settings*

- *Covers a few different areas and helpful in clinical trials*
- *Helpful to differentiate MCI and AD*
- *May have a good correlation with PET and CSF measures*
- *Not used in the clinical setting and limited clinical relevance*

*This is the experience and thoughts of our expert authors



Disability Assessment for Dementia (DAD)

The DAD is a clinical and functional observer-reported outcome measure used to assess the ability to perform activities, tasks, and behaviors of everyday life in people with dementia.⁴¹ Activities are organized such that the ability to initiate, plan, and execute each is evaluated, therefore, informing on cognitive ability as well. Caregivers or other informants complete the questionnaire. It is designed to be free of gender bias and is performance-based.⁴¹

40 items on basic self-care, instrumental ADLs, and leisure:⁴¹

Basic ADLs (activities important for self-care), including dressing, hygiene, continence, and eating

Instrumental ADLs (activities important for maintenance in a specific environment), including meal preparation, telephoning, housework, taking care of finance and correspondence, going on an outing, taking medications, and the ability to stay safely at home

Leisure activities (activities beyond self-maintenance that are for the purpose of recreation), in terms of the interest that is shown toward these activities

A higher global score represents less disability

Insights on use in clinical practice and research settings*

- *Reflects patient outcomes in relation to day-to-day function*
- *Some parts of the test are used by occupational therapists in memory clinics*
- *Not reflecting the clinical trials at early stages*

*This is the experience and thoughts of our expert authors



Functional Activities Questionnaire (FAQ)

The FAQ is an observer-reported outcome used across many diseases. It is a clinical and functional measure of instrumental activities of daily living as assessed by an informant (e.g. a caregiver or family member) over 4 weeks.⁴² A total FAQ score ≥ 6 may be used to distinguish mild AD from MCI.⁴² The FAQ has been shown to predict progression from MCI to AD.⁷³ However, if the patient has not performed a certain activity, the questionnaire will be incomplete.⁴²

Instrumental ADLs assessed on the FAQ:⁴²

<ul style="list-style-type: none"> • Writing checks, paying bills, and keeping financial records 	<ul style="list-style-type: none"> • Keeping track of current events
<ul style="list-style-type: none"> • Assembling tax or business records 	<ul style="list-style-type: none"> • Attending to and understanding a television program, book, or magazine
<ul style="list-style-type: none"> • Shopping alone 	<ul style="list-style-type: none"> • Remembering appointments, family occasions, and medications
<ul style="list-style-type: none"> • Playing a game of skill 	<ul style="list-style-type: none"> • Traveling out of the neighborhood
<ul style="list-style-type: none"> • Making coffee or tea 	<ul style="list-style-type: none"> • Preparing a balanced meal

Performance in each category is rated as:⁴²

- 0 = *normal*
- 1 = *has difficulty, but does by self*
- 2 = *requires assistance*
- 3 = *dependent*

Score ranges from 0 to 30; higher scores on each of the categories indicate greater functional impairment⁴²

Insights on use in clinical practice and research settings*

- *Used in some clinics, mainly by occupational therapists*
- *Shows patient decline as a set of standards better than general questioning*
- *Highly translational and measures of ADL provide meaningful measures and pragmatic assessment of clinical change*

*This is the experience and thoughts of our expert authors



Amsterdam - Instrumental Activities of Daily Living (A-IADL)

The A-IADL is a 70-item observer-reported questionnaire aimed at detecting early dementia and early-onset dementia.⁴³ It is a registered trademark of Alzheimer Center VU University Medical Center (Amsterdam, The Netherlands).^{43,74} There is also a short version, with 30 items, which still covers the range of functional domains and takes 10–15 minutes to complete.⁷⁵ An informant of the patient (e.g. a caregiver or family member) completes the questionnaire.^{43,74} IADL are complex activities that require multiple cognitive processes and involve minimal automated skills.⁴³

70 items or 30 items (short-version), pertaining to, e.g. the following:^{43,75}

• Household duties	• Paperwork and computer use
• Shopping	• Operating devices
• Cooking	• Driving and transportation
• Domestic appliances	• Games and books
• Finances and paying bills	

Each item has a 5-point scale response option:⁴³

- 0 = no difficulty
- 1 = slightly more difficulty
- 2 = more difficulty
- 3 = much more difficulty
- 4 = no longer able to perform this task

Higher scores indicate poorer functioning⁷⁴

Insights on use in clinical practice and research settings*

- *Becoming a preferred measure in trials due to its sensitivity and contemporary activities/questions*
- *Not used in clinical practice*

*This is the experience and thoughts of our expert authors



Quality of Life in Alzheimer's Disease (QoL-AD)

The QoL-AD assesses quality of life, physical condition, mood, interpersonal relationships, ability to participate in meaningful activity, and financial situation.^{4,46} Both self-assessment and informant interview versions exist; each can be completed in under 10 minutes.⁶ Proxy-rated QoL-AD is typically found to be lower than self-rated QoL-AD;^{47,48} and self-assessed QoL-AD may have limited correlation with changes in clinical outcomes.⁴⁷

Domains:⁴⁷

• Physical health	• Friends
• Energy	• Self as a whole
• Mood	• Ability to do chores around the house
• Living situation	• Ability to do things for fun
• Memory	• Money
• Family	• Life as a whole
• Marriage	

Each domain is assessed on a 4-point scale:^{46,48}

- 1 = *poor*
- 2 = *fair*
- 3 = *good*
- 4 = *excellent*

Total score ranges from 13 to 52; a higher score represents better quality of life⁴⁶

Insights on use in clinical practice and research settings*

- *Wide range of scoring/testing is helpful for clinical trials*
- *Short and quick*
- *Not routinely used in clinical care*

*This is the experience and thoughts of our expert authors



5-Level EuroQoL-5D (EQ-5D-5L)

The EQ-5D-5L is a proxy-rated outcome measure used across many diseases to measure the immediate health of the patient, i.e. “health today”.^{49,50,76} It is designed for self-completion but may be used by a caregiver or family member instead. As a self-completed measure, it may not accurately reflect clinically important changes.⁵⁰

The **EQ-5D-5L** assesses **5 dimensions**:⁴⁹

- *Mobility*
- *Self-care*
- *Usual activities*
- *Pain/discomfort*
- *Anxiety/depression*

Each dimension has 5 response categories:⁷⁶

- *No problems*
- *Slight problems*
- *Moderate problems*
- *Severe problems*
- *Unable to complete task*

An index score is calculated from -0.281 to 1 ⁷⁶

Higher scores indicate better quality of life⁷⁶

Insights on use in clinical practice and research settings*

- *Used mostly in the research setting*

*This is the experience and thoughts of our expert authors



Alzheimer's Disease Composite Score (ADCOMS)

The ADCOMS is a composite cognitive and functional endpoint that combines elements of 3 common endpoints (CDR-SB, ADAS-Cog, and MMSE) with the total score ranging from 0 to 1.97.²⁴ As such it is both a performance outcome and an observer-reported outcome. ADCOMS is intended for use in the clinical research setting. Sensitivity to clinical change in MCI/prodromal AD has been reported to be improved with the ADCOMS compared with the CDR-SB, ADAS-Cog, and MMSE individually.²⁴

Scale and item ²⁴	PLS coefficient (weight)
ADAS-Cog	
Delayed word recall	0.008
Orientation	0.017
Word recognition	0.004
Word finding difficulty	0.016
MMSE	
Orientation time	0.042
Drawing	0.038
CDR-SB	
Personal care	0.054
Community affairs	0.109
Home and hobbies	0.089
Judgement and problem solving	0.069
Memory	0.059
Orientation	0.078

Insights on use in clinical practice and research settings*

- *Not used in everyday practice as measures must be statistically combined*
- *ADAS-Cog (the key component) is not commonly done in clinical practice*
- *Heavily weighted towards orientation (from all 3 tests)*

*This is the experience and thoughts of our expert authors



Cognitive Function Instrument (CFI)

The CFI is an assessment of early changes in cognitive status and functional abilities.²⁵ The CFI is a questionnaire completed by patients and/or an informant (e.g. a caregiver or family member).^{26,27} Examples of questions are:²⁶ Do others tell you that you tend to repeat questions over and over? And do you need more help from others to remember appointments, family occasions, or holidays? The index was originally developed to determine whether a self-assessed measure of change in cognition is reliable and may serve as a trial endpoint.²⁵

14 questions including items on the following (over the previous year):^{26,27}

- *Memory decline*
- *Cognitive difficulties*
- *Functional decline*

Responses are recorded as:^{26,27}

- *Yes = 1*
- *No = 0*
- *Maybe = 0.5*
- *Not applicable (an option for questions regarding driving, handling finances, and word performance)*

Responses are summed to create a total score, ranging from 0 to 14; a higher score reflects greater impairment^{26,27}

Insights on use in clinical practice and research settings*

- *Helpful as a self-assessment for patient outcome measures in primary care*
- *Sensitive in early disease*
- *Minimal use in the clinical setting*
- *Lacks the sensitivity required for clinical trials*

*This is the experience and thoughts of our expert authors



Integrated Alzheimer's Disease Rating Scale (iADRS)

The iADRS combines scores from the ADAS-Cog and the ADCS-iADL.²⁹ Scores range from 0 to 146 with higher scores indicating better performance.²⁹ This composite score was designed to be more sensitive to MCI and to reduce variability in data.²⁹ Its sensitivity to progression from MCI to moderate AD has been reported as superior to several other composites and equal to or better than individual scales in detecting treatment differences.²⁸

The iADRS can be divided into 2 principal components: cognitive items and instrumental ADL.²⁸

Items:²⁸

Cognition

Word recognition
 Delayed word recall
 Word recall
 Orientation

Daily function

Making a meal
 Telephone
 Going shopping
 Household appliance
 Current events
 Obtaining beverage
 Writing
 Getting around
 Being alone
 Television
 Keeping appointments
 Clearing dishes

Insights on use in clinical practice and research settings*

- *Rarely used in clinical practice*
- *Degree to which it reflects clinical trial results and patient outcomes is not yet known*

*This is the experience and thoughts of our expert authors



Alzheimer's Disease Cooperative Study - Clinical Global Impression of Change (ADCS-CGIC)

The ADCS-CGIC is designed to assess clinically meaningful change over time in cognition, function, and behavior, in the research setting.^{30,31} Unlike a targeted symptom scale, ADCS-CGIC takes into account the patient's overall function in these 3 domains.³⁰ The ADCS-CGIC is completed by a clinician who performs direct interviews with both the patient and the informant/caregiver.³¹

Domains:^{30,31}

- *General*
- *Mental cognitive state*
- *Activities of daily living*
- *Behavior*

ADCS-CGIC uses a **7-point** scale:^{30,31}

- *1 = marked improvement*
- *2 = moderate improvement*
- *3 = minimal improvement*
- *4 = no change*
- *5 = minimal worsening*
- *6 = moderate worsening*
- *7 = marked worsening*

Insights on use in clinical practice and research settings*

- *Not used in clinical practice, but easily understandable, meaningful endpoint*
- *Reflects patient outcomes, as it incorporates both patient and caregiver interview performed (and interpreted) by physician*
- *Used in clinical trials in several disease stages, from preclinical to mild AD*

*This is the experience and thoughts of our expert authors



Neuropsychiatric Inventory Questionnaire (NPI-Q)

The NPI is an interview of the caregiver or other informant assessing neuropsychiatric symptoms in the past 4 weeks, across 12 domains.⁵¹ The NPI-Q is a condensed version of the NPI, completed by the caregiver/informant.⁵¹ Symptom severity is assessed, but unlike the NPI, frequency is not assessed. Neuropsychiatric symptom severity on the NPI-Q has been shown to be associated with cognitive and functional impairment.⁷⁷ The NPI-Q is intended for use in clinical practice and is typically completed in under 5 minutes.⁵¹

Domains:⁵¹

• Delusions	• Disinhibition
• Hallucinations	• Irritability/lability
• Agitation/aggression	• Aberrant motor behavior
• Dysphoria/depression	• Nighttime disturbances
• Euphoria/elation	• Appetite/eating disturbances
• Apathy/indifference	

Each domain is assessed for severity on a 3-point scale:

- 1 = *mild*
- 2 = *moderate*
- 3 = *severe*

Total severity score ranges from 0 to 36; a higher score represents greater symptom burden

Insights on use in clinical practice and research settings*

- *Occasionally used in the clinical setting*
- *Considered useful in clinical practice and clinical trials for behavioural and psychological symptoms of dementia*

*This is the experience and thoughts of our expert authors



Columbia Suicide Severity Rating Scale (C-SSRS)

The C-SSRS is a safety outcome used across the field of psychiatry to prospectively measure the severity and frequency of suicidal ideation and suicidal behavior.^{52,53} It is an observer-reported outcome in the form of a clinical interview intended to aid clinicians in assessing patient suicide risk.⁵³ Four constructs of suicide ideation and behavior are measured.⁵² However, it should be noted that the C-SSRS is not specifically validated in the elderly population;⁵³ and concerns have been raised regarding the appropriateness of the C-SSRS for use in patients with dementia.⁵⁴

C-SSRS components:⁵²

- **Severity of ideation** (yes/no):
 1. *Wish to be dead*
 2. *Non-specific active suicidal thoughts*
 3. *Suicidal thoughts with methods*
 4. *Suicidal intent*
 5. *Suicidal intent with plan*
- **Intensity of ideation:**
 - *Frequency, duration, controllability, deterrents, and reason for ideation*
 - *Each rated on a 5-point scale (1 being the least severe and 5 being the most severe)*
- **Behavior subscale** (yes/no):
 - *Actual, aborted, and interrupted attempts; preparatory behavior; and non-suicidal self-injurious behavior*
- **Lethality subscale:**
 - *Actual lethality/medical damage*
 - *6-point scale (0 being no/minor physical damage and 5 being death)*
 - *If actual lethality is 0, potential lethality of attempts is assessed*
 - *3-point ordinal scale (0 being least likely lethal and 2 being most likely lethal)*

Insights on use in clinical practice and research settings*

- *Important safety assessment that can be done by trained raters, including non-experts in behavioral health; therefore, more useful in daily practice*
- *A safety measure in clinical trials but not used in clinical settings*
- *Does not adequately reflect patient outcomes and is inferior to a standard risk assessment*

*This is the experience and thoughts of our expert authors



Zarit Burden Interview (ZBI)

The ZBI evaluates caregiver burden in dementia.⁵⁵ The following domains are addressed through a series of statements: caregiver health, psychological well-being, relationship with the patient, social life, and finances.⁵⁵

The original **ZBI** has 29 items⁵⁵

Examples of items assessed:

- *I feel embarrassed over my spouse's behavior*
- *I feel strained in my interactions with my spouse*
- *I feel that my health has suffered because of my involvement with my spouse*
- *I feel that my spouse doesn't appreciate what I do for him/her as much as I would like*

Insights on use in clinical practice and research settings*

- *Used in some clinics and useful as a secondary outcome in clinical trials*
- *Reflects caregiver burden but not directly related to severity of patient's disease*

*This is the experience and thoughts of our expert authors



Resource Utilization in Dementia (RUD)

The RUD, developed for use in cost-effectiveness studies, captures resource use for dementia care.⁵⁶ The tool was designed for multinational use and to capture costs from the societal perspective. Both formal and informal care are assessed.⁵⁶ The RUD-Lite is an abbreviated version of the RUD and focuses on patient resource use (rather than both patient and caregiver resource use).⁵⁷

The **RUD** tool assesses:^{56,57}

- *Time spent assisting the patient with ADLs and instrumental ADLs*
- *Time spent supervising the patient*
- *Patient living accommodations*
- *Patient healthcare resource utilization*
- *Caregiver work status/impact, days missed*
- *Caregiver sleep*
- *Caregiver healthcare resource utilization*

Insights on use in clinical practice and research settings

- *Not used in the clinical setting, used in some service evaluation programs*
- *Helpful in clinical trials but may not necessarily reflect resource use during the trial itself; resource use tends to be delayed*

*This is the experience and thoughts of our expert authors



Fluid-based and imaging biomarkers in AD clinical trials

The FDA-NIH Biomarker Working Group defines a biomarker as:⁷⁸ “A defined characteristic that is measured as an indicator of normal biological processes, pathogenic processes, or responses to an exposure or intervention, including therapeutic interventions. Molecular, histologic, radiographic, or physiologic characteristics are types of biomarkers. A biomarker is not an assessment of how an individual feels, functions, or survives.”

In AD clinical trials, biomarkers can be classified as either diagnostic or endpoint biomarkers.⁷⁹

- **Diagnostic biomarkers** may be used for inclusion/exclusion criteria, sample size determination, or increasing statistical power (e.g. via stratification)
- **Endpoint biomarkers**, as trial outcomes, measure disease progression and detect treatment effects


The evolving characterization of AD as a clinico-biological diagnosis requires biomarkers that detect preclinical and prodromal AD.⁸⁰ In addition to the early detection of AD pathophysiology for the inclusion of patients in clinical trials, biomarkers may be used for enrichment of the patient population by identifying those potentially most likely to experience treatment benefit.⁸⁰ Amyloid positron emission tomography (PET) and tau PET tracers are increasingly used as endpoints in AD clinical trials and play an important role in this setting.⁸¹

AD biomarkers may be biochemical- or imaging-based. Biochemical markers such as A β 42, total Tau (tTau), phosphorylated Tau (pTau), A β 42:40 ratio, and pTau:A β 42 ratio are quantitative. Imaging biomarkers are qualitative (continuous) or quantitative (positive/negative) and include structural imaging such as magnetic resonance imaging (MRI) or computed tomography (CT); functional imaging such as fluorodeoxyglucose (FDG)-PET, functional MRI (fMRI), or single photon emission computed tomography (SPECT); or molecular imaging such as amyloid-PET or Tau-PET.

Biomarkers are essential for the detection of AD and the measurement of biological progression, and assays exist to measure both amyloid and tau progression (**Figure 5**).^{82–88} Elecsys[®] cerebrospinal fluid (CSF) assays are commercially available in CE Mark accepting countries (28 EU countries plus Iceland, Lichtenstein, Norway, and Switzerland) and have FDA breakthrough device designation.⁸⁵ The assays are intended for use in adult subjects with cognitive impairment or MCI being evaluated for AD, and as an adjunct to their clinical diagnostic evaluations.

Figure 5: Assays that detect amyloid and tau progression

Note: the figure below shows Roche assays and is not intended to be comprehensive.



*Elecsys[®] CSF assays are not approved or cleared for clinical use in the US.

Aβ, amyloid beta; AD, Alzheimer’s Disease; CSF, cerebrospinal fluid; FDA, Food and Drug Administration; MCI, mild cognitive impairment.

Individual biomarkers summarized

In this section, we summarize biomarkers used in AD, including the scientific rationale for their use in clinical trials to diagnose MCI and AD.

CSF – amyloid beta (A β)

Low CSF levels of A β 42 is a key feature of AD.⁸⁹ A β is sequestered into amyloid plaques, reducing the amount in the CSF, and A β is predictive of AD pathology – CSF A β 42 is highly sensitive for identification of prodromal AD cases in MCI cohorts.^{89,90}

Insights on use in clinical practice and research settings*

- *Use in clinical practice depends on cost (of test), reimbursement rates (for lumbar puncture), and regional differences in patient willingness to undergo procedure*
- *Large variability in use depending on practice patterns in that area / expertise*
- *Strongly reflects clinical trial outcomes and key features of AD; pre vs post change could be useful*
- *Reflection of patient outcomes is likely delayed*

*This is the experience and thoughts of our expert authors



CSF – tau

High levels of pTau are characteristic of prodromal AD and AD dementia.^{89,91,92} pTau is a measure of the amount of tau that is phosphorylated, the variant of tau found in tangles.⁹¹ pTau predicts progression from MCI to AD dementia.⁹²

Increased levels of tTau is characteristic of prodromal AD and AD dementia, reflecting axonal (neuronal) degeneration.^{89,91} tTau gives a measure of the intensity of neurodegeneration in AD, but it is not a disease-specific marker.⁸⁹

Insights on use in clinical practice and research settings*

- *Use in clinical practice depends on cost (of test), reimbursement rates (for lumbar puncture), and regional differences in patient willingness to undergo procedure*
- *Large variability in use depending on practice patterns in that area / expertise*
- *Strongly reflects clinical trial outcomes and key features of AD; pre vs post change could be useful*
- *As it is measurable, it can show the level of decline, but it must be combined with amyloid biomarkers*
- *Reflective of patient outcomes (assuming correct diagnosis of AD)*

*This is the experience and thoughts of our expert authors



CSF – amyloid beta and tau ratios

A β ratios are predictive of AD pathology.^{89,91} a decreased A β 42:A β 40 ratio in MCI is predictive of progression to dementia.⁸⁹ A β 40 is the most prevalent form of A β in CSF – the A β 42:40 ratio accounts for between-individual differences in A β isoforms.⁹¹

High tTau:A β 42 and pTau:A β 42 ratios predict future cognitive decline.^{93,94} A combined CSF biomarker test using A β 42 and tTau has been reported to have 95% sensitivity and 83% specificity for the prediction of progression to AD.⁹³

In the BioFINDER cohort, Elecsys[®] CSF assays were used to analyze a variety of CSF biomarkers.⁸⁴ pTau/A β 42 and tTau/A β 42 had \approx 90% concordance with amyloid PET imaging, outperforming A β 42 concordance (80%). Specificity and area under the curve were also improved with pTau/A β 42 and tTau/A β 42 ratios compared with A β 42.⁸⁴

Insights on use in clinical practice and research settings*

- *Ratios add substantial value for clinical trials when obtaining longitudinal assessments; these are good clinical outcome measures*
- *Use in the clinic is increasing but still in early stages of adoption*

*This is the experience and thoughts of our expert authors



Amyloid PET

Amyloid tracer binds to fibrillar A β and amyloid retention can be estimated by PET signal.⁹⁵ Types of tracers include Pittsburgh compound-B, florbetapir [18F] [AmyvidTM], florbetaben [18F] [NeuraceqTM], and flutemetamol [18F] [VizamylTM]. There are 3 FDA-approved tracers.

Insights on use in clinical practice and research settings*

- *Mixed opinion on utility for clinical trials: thought to be useful for baseline and longitudinal measures but also noted to be difficult to measure change*
- *In early stages of adoption in clinical practice*
- *Whether reflective of patient outcomes is a key question but believed by some to reflect outcomes in most cases*
- *Cost and exposure to radiation are potential limitations*

*This is the experience and thoughts of our expert authors



Tau PET

Tau PET tracers have been designed to bind neurofibrillary tangles, tau aggregates, and neuropil threads.⁹⁶ Flortaucipir [TauvidTM] is the only ligand which is FDA approved for tau PET in those with suspected AD.^{97,98} [18F]GTP1 is a novel tau PET tracer in development that binds to tau pathology, enabling study of tau propagation.^{86,99} GTP1 detects change in tau pathology over time in untreated patients.¹⁰⁰ Uptake of [18F]GTP1 has also been shown to correlate with the degree of cognitive impairment in AD.^{101,102}

Insights on use in clinical practice and research settings*

- *In early stages of adoption in clinical practice; use in clinical trials and whether reflective of patient outcomes are to be determined*
- *Likely helpful for optimal timing of targeted therapies: anti-amyloid (e.g. low to moderate amyloid and no or minimal tau), and for anti-tau therapies and anti-neuroinflammation therapies*

*This is the experience and thoughts of our expert authors



Volumetric MRI

Structural brain changes that occur with AD can be observed even prior to appearance of clinical symptoms.¹⁰³ Hippocampal volume is reduced prior to AD diagnosis. In dementia due to AD, entorhinal cortex volume is decreased by $\approx 30\%$ – 40% and hippocampal volume by $\approx 15\%$ – 25% compared with controls.¹⁰⁴ Structural features on MRI can predict progression from MCI to AD.¹⁰⁵

Insights on use in clinical practice and research settings*

- *Widely used in clinical practice, and often underutilized by both neurologists and primary care physicians*
- *Hippocampal atrophy in setting of progressive short-term memory loss and family history of AD is highly predictive and cost-effective*
- *Radiologists can be asked to comment on hippocampal and other regional atrophy, saving time for other clinicians*
- *Often used in trials at baseline and post-intervention; reflective of clinical trial results but with a 6–12-month delay compared with CSF biomarkers*
- *Not linearly/directly reflective of patient outcomes*

*This is the experience and thoughts of our expert authors



Potential digital biomarkers

Prior to the onset of AD symptoms, biomarkers have the potential to predict AD progression.¹⁰⁶ Changes in olfaction, hearing, and walking speed can occur 5–15 years prior to the onset of cognitive impairments, and are therefore strong indicators of dementia.¹⁰⁷ Sensory and motor manifestations of AD can occur up to 15 years prior to an effective diagnosis; however, cognitive tests take a long time to administer, are limited by cultural bias, and are rater dependent.¹⁰⁸ In addition, stigma around AD may limit the use of cognitive testing and/or lead to delayed diagnosis.^{109,110}

Digital biomarkers may have the potential to accelerate AD diagnosis and may improve prognosis.¹⁰⁸ Adoption of increasingly sophisticated mobile and wearable technologies (e.g. smart phones and smart watches) offers the opportunity to use “digital biomarkers” to measure the early changes in sensory and motor signs of AD. These measures can be passive/acquired (heart rate variability; gait speed; GPS) or active/prompted (memory tests; tapping tests; voice tests; eye movements).¹⁰⁸

Summary

- *A variety of endpoints have been developed and used in AD clinical trials to assess potential therapeutic effect*
- *These outcome measures include cognitive, functional, quality of life, psychiatric, and caregiver burden scales, as well as imaging and biomarkers*
- *Utility of individual endpoints in clinical practice varies considerably, as does the extent to which they reflect clinical trial results and patient outcomes*

References

1. FDA. Patient-Focused Drug Development Guidance Public Workshop: methods to identify what is important to patients & select, develop or modify fit-for-purpose clinical outcomes assessments. Appendix. <https://www.fda.gov/media/116281/download>. Accessed March 4, 2021.
2. FDA. Patient-Focused Drug Development: methods to identify what is important to patients. <https://www.fda.gov/media/131230/download>. Accessed March 4, 2021.
3. FDA. Multiple endpoints in clinical trials guidance for industry. 2018 <https://www.fda.gov/media/102657/download>. Accessed February 2021.
4. Logsdon RG, Gibbons LE, McCurry SM, Teri L. Quality of life in Alzheimer's disease: patient and caregiver reports. *J Ment Health Aging*. 1999;5:21–32.
5. Benoit JS, Chan W, Piller L, Doody R. Longitudinal sensitivity of Alzheimer's disease severity staging. *Am J Alzheimers Dis Other Demen*. 2020;35:1533317520918719.
6. EMA. Guideline on the clinical investigation of medicines for the treatment of Alzheimer's disease. 2018 https://www.ema.europa.eu/en/documents/scientific-guideline/guideline-clinical-investigation-medicines-treatment-alzheimers-disease-revision-2_en.pdf. Accessed February 2021.
7. Walton M. Endpoints in alzheimer's disease clinical trials. ISCTM Annual Meeting 2016.
8. Webster L, Groskreutz D, Grinbergs-Saull A, Howard R, O'Brien JT, Mountain G, Banerjee S, Woods B, Pernecky R, Lafortune L, Roberts C, McCleery J, Pickett J, Bunn F, Challis D, Charlesworth G, Featherstone K, Fox C, Goodman C, Jones R, Lamb S, Moniz-Cook E, Schneider J, Shepperd S, Surr C, Thompson-Coon J, Ballard C, Brayne C, Burns A, Clare L, Garrard P, Kehoe P, Passmore P, Holmes C, Maidment I, Robinson L, Livingston G. Core outcome measures for interventions to prevent or slow the progress of dementia for people living with mild to moderate dementia: systematic review and consensus recommendations. *PLoS One*. 2017;12:e0179521.
9. FDA. Early Alzheimer's Disease: developing drugs for treatment guidance for industry. 2018 <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/alzheimers-disease-developing-drugs-treatment-guidance-industry>. Accessed February 2021.
10. Schneider LS, Goldberg TE. Composite cognitive and functional measures for early stage Alzheimer's disease trials. *Alzheimers Dement (Amst)*. 2020;12:e12017.
11. Weintraub S, Carrillo MC, Farias ST, Goldberg TE, Hendrix JA, Jaeger J, Knopman DS, Langbaum JB, Park DC, Ropacki MT, Sikkes SAM, Welsh-Bohmer KA, Bain LJ, Brashear R, Budur K, Graf A, Martenyi F, Storck MS, Randolph C. Measuring cognition and function in the preclinical stage of Alzheimer's disease. *Alzheimers Dement (N Y)*. 2018;4:64–75.
12. ClinicalTrials.gov. U.S. National Library of Medicine. NCT03444870 <https://clinicaltrials.gov/ct2/show/NCT03444870>. Accessed February 2021.
13. ClinicalTrials.gov. U.S. National Library of Medicine. NCT03443973. <https://clinicaltrials.gov/ct2/show/NCT03443973>. Accessed February 2021.
14. ClinicalTrials.gov. U.S. National Library of Medicine. NCT03828747. <https://clinicaltrials.gov/ct2/show/NCT03828747>. Accessed February 2021.
15. ClinicalTrials.gov. U.S. National Library of Medicine. NCT03289143. <https://clinicaltrials.gov/ct2/show/NCT03289143>. Accessed February 2021.
16. ClinicalTrials.gov. U.S. National Library of Medicine. NCT02484547. <https://clinicaltrials.gov/ct2/show/NCT02484547>. Accessed February 2021.
17. ClinicalTrials.gov. U.S. National Library of Medicine. NCT02477800. <https://clinicaltrials.gov/ct2/show/NCT02477800>. Accessed February 2021.
18. ClinicalTrials.gov. U.S. National Library of Medicine. NCT04468659. <https://clinicaltrials.gov/ct2/show/NCT04468659>. Accessed February 2021.
19. ClinicalTrials.gov. U.S. National Library of Medicine. NCT03887455. <https://clinicaltrials.gov/ct2/show/NCT03887455>. Accessed February 2021.
20. ALZ-801: Alzheon pipeline. <https://alzheon.com/pipeline/alzheon-alz-801/>. Accessed February 2021.
21. ClinicalTrials.gov. U.S. National Library of Medicine. NCT04437511. <https://clinicaltrials.gov/ct2/show/NCT04437511>. Accessed February 2021.
22. Morris JC. The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology*. 1993;43:2412–2414.
23. Cedarbaum JM, Jaros M, Hernandez C, Coley N, Andrieu S, Grundman M, Vellas B. Rationale for use of the Clinical Dementia Rating Sum of Boxes as a primary outcome measure for Alzheimer's disease clinical trials. *Alzheimers Dement*. 2013;9(1 Suppl):S45–S55.
24. Wang J, Logovinsky V, Hendrix SB, Stanworth SH, Perdomo C, Xu L, Dhadda S, Do I, Rabe M, Luthman J, Cummings J, Satlin A. ADCOMS: a composite clinical outcome for prodromal Alzheimer's disease trials. *J Neurol Neurosurg Psychiatry*. 2016;87:993–999.




25. Li C, Neugroschl J, Luo X, Zhu C, Aisen P, Ferris S, Sano M. The utility of the Cognitive Function Instrument (CFI) to detect cognitive decline in non-demented older adults. *J Alzheimers Dis.* 2017;60:427–437.
26. Walsh SP, Raman R, Jones KB, Aisen PS. ADCS Prevention Instrument Project: the Mail-In Cognitive Function Screening Instrument (MCFSI). *Alzheimer Dis Assoc Disord.* 2006;20(Suppl 3):S170–S178.
27. Amariglio RE, Donohue MC, Marshall GA, Rentz DM, Salmon DP, Ferris SH, Karantzoulis S, Aisen PS, Sperling RA. Tracking early decline in cognitive function in older individuals at risk for Alzheimer disease dementia: the Alzheimer's Disease Cooperative Study Cognitive Function Instrument. *JAMA Neurol.* 2015;72:446–454.
28. Wessels AM, Siemers ER, Yu P, Andersen SW, Holdridge KC, Sims JR, Sundell K, Stern Y, Rentz DM, Dubois B, Jones RW, Cummings J, Aisen PS. A combined measure of cognition and function for clinical trials: the Integrated Alzheimer's Disease Rating Scale (iADRS). *J Prev Alzheimers Dis.* 2015;2:227–241.
29. Wessels AM, Andersen SW, Dowsett SA, Siemers ER. The Integrated Alzheimer's Disease Rating Scale (iADRS) findings from the EXPEDITION3 trial. *J Prev Alzheimers Dis.* 2018;5:134–136.
30. Schneider LS, Clark CM, Doody R, Ferris SH, Morris JC, Raman R, Reisberg B, Schmitt FA. ADCS Prevention Instrument Project: ADCS-clinicians' global impression of change scales (ADCS-CGIC), self-rated and study partner-rated versions. *Alzheimer Dis Assoc Disord.* 2006;20(Suppl 3):S124–S138.
31. Schneider LS, Olin JT, Doody RS, Clark CM, Morris JC, Reisberg B, Schmitt FA, Grundman M, Thomas RG, Ferris SH. Validity and reliability of the Alzheimer's Disease Cooperative Study-Clinical Global Impression of Change. The Alzheimer's Disease Cooperative Study. *Alzheimer Dis Assoc Disord.* 1997;11(Suppl 2):S22–S32.
32. Li K, Chan W, Doody RS, Quinn J, Luo S. Prediction of conversion to Alzheimer's disease with longitudinal measures and time-to-event data. *J Alzheimers Dis.* 2017;58:361–371.
33. Rosen WG, Mohs RC, Davis KL. A new rating scale for Alzheimer's disease. *Am J Psychiatry.* 1984;141:1356–1364.
34. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975;12:189–198.
35. Arevalo-Rodriguez I, Smailagic N, Roqué IFM, Ciapponi A, Sanchez-Perez E, Giannakou A, Pedraza OL, Bonfill Cosp X, Cullum S. Mini-Mental State Examination (MMSE) for the detection of Alzheimer's disease and other dementias in people with mild cognitive impairment (MCI). *Cochrane Database Syst Rev.* 2015;2015:CD010783.
36. Sutin AR, Stephan Y, Terracciano A. Verbal fluency and risk of dementia. *Int J Geriatr Psychiatry.* 2019;34:863–867.
37. Weiss LG, Saklofske DH, Coalson D, Raiford SE, eds. *WAIS-IV clinical use and interpretation: Scientist-practitioner perspectives.* 1st ed. Academic Press; 2010.
38. Mura T, Proust-Lima C, Jacqmin-Gadda H, Akbaraly TN, Touchon J, Dubois B, Berr C. Measuring cognitive change in subjects with prodromal Alzheimer's disease. *J Neurol Neurosurg Psychiatry.* 2014;85:363–370.
39. Randolph C, Tierney MC, Mohr E, Chase TN. The Repeatable Battery for the Assessment of Neuropsychological Status (RBANS): preliminary clinical validity. *J Clin Exp Neuropsychol.* 1998;20:310–319.
40. Donohue MC, Sperling RA, Salmon DP, Rentz DM, Raman R, Thomas RG, Weiner M, Aisen PS. The preclinical Alzheimer cognitive composite: measuring amyloid-related decline. *JAMA Neurol.* 2014;71:961–970.
41. Gélinas I, Gauthier L, McIntyre M, Gauthier S. Development of a functional measure for persons with Alzheimer's disease: the disability assessment for dementia. *Am J Occup Ther.* 1999;53:471–481.
42. Teng E, Becker BW, Woo E, Knopman DS, Cummings JL, Lu PH. Utility of the functional activities questionnaire for distinguishing mild cognitive impairment from very mild Alzheimer disease. *Alzheimer Dis Assoc Disord.* 2010;24:348–353.
43. Sikkes SA, de Lange-de Klerk ES, Pijnenburg YA, Gillissen F, Romkes R, Knol DL, Uitdehaag BM, Scheltens P. A new informant-based questionnaire for instrumental activities of daily living in dementia. *Alzheimers Dement.* 2012;8:536–543.
44. Galasko D, Bennett D, Sano M, Ernesto C, Thomas R, Grundman M, Ferris S. An inventory to assess activities of daily living for clinical trials in Alzheimer's disease. The Alzheimer's Disease Cooperative Study. *Alzheimer Dis Assoc Disord.* 1997;11(Suppl 2):S33–S39.
45. Goldberg TE, Koppel J, Keehlisen L, Christen E, Dreses-Werringloer U, Conejero-Goldberg C, Gordon ML, Davies P. Performance-based measures of everyday function in mild cognitive impairment. *Am J Psychiatry.* 2010;167:845–853.
46. Logsdon RG, Gibbons LE, McCurry SM, Teri L. Assessing quality of life in older adults with cognitive impairment. *Psychosom Med.* 2002;64:510–519.
47. Lacey L, Bobula J, Rudell K, Alvir J, Leibman C. Quality of life and utility measurement in a large clinical trial sample of patients with mild to moderate Alzheimer's disease: determinants and level of changes observed. *Value Health.* 2015;18:638–645.
48. Romhild J, Fleischer S, Meyer G, Stephan A, Zwakhalen S, Leino-Kilpi H, Zabalegui A, Saks K, Soto-Martin M, Sutcliffe C, Rahm Hallberg I, Berg A, RightTimePlaceCare C. Inter-rater agreement of the Quality of Life-Alzheimer's Disease (QoL-AD) self-rating and proxy rating scale: secondary analysis of RightTimePlaceCare data. *Health Qual Life Outcomes.* 2018;16:131.
49. Herdman M, Gudex C, Lloyd A, Janssen M, Kind P, Parkin D, Bonsel G, Badia X. Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res.* 2011;20:1727–1736.
50. Martin A, Meads D, Griffiths AW, Surr CA. How should we capture health state utility in dementia? Comparisons of DEMQOL-Proxy-U and of self- and proxy-completed EQ-5D-5L. *Value Health.* 2019;22:1417–1426.

51. Kaufer DI, Cummings JL, Ketchel P, Smith V, MacMillan A, Shelley T, Lopez OL, DeKosky ST. Validation of the NPI-Q, a brief clinical form of the Neuropsychiatric Inventory. *J Neuropsychiatry Clin Neurosci*. 2000;12:233–239.
52. Posner K, Brown GK, Stanley B, Brent DA, Yershova KV, Oquendo MA, Currier GW, Melvin GA, Greenhill L, Shen S, Mann JJ. The Columbia–Suicide Severity Rating Scale: initial validity and internal consistency findings from three multisite studies with adolescents and adults. *Am J Psychiatry*. 2011;168:1266–1277.
53. Alphas L, Brashear HR, Chappell P, Conwell Y, Dubrava S, Khin NA, Kozauer N, Hartley DM, Miller DS, Schindler RJ, Siemers ER, Stewart M, Yaffe K. Considerations for the assessment of suicidal ideation and behavior in older adults with cognitive decline and dementia. *Alzheimers Dement (N Y)*. 2016;2:48–59.
54. Chappell P, Dubrava S, Stewart M, Hartley DM, Alphas L, Brashear HR, Conwell Y, Miller D, Schindler RJ, Siemers ER, Yaffe K. Suicidal ideation and behavior assessment in dementia studies: an internet survey. *Alzheimers Dement (N Y)*. 2016;2:60–68.
55. Zarit SH, Reever KE, Bach-Peterson J. Relatives of the impaired elderly: correlates of feelings of burden. *Gerontologist*. 1980;20:649–655.
56. Wimo A, Jonsson L, Zbrozek A. The Resource Utilization in Dementia (RUD) instrument is valid for assessing informal care time in community-living patients with dementia. *J Nutr Health Aging*. 2010;14:685–690.
57. Wimo A, Gustavsson A, Jönsson L, Winblad B, Hsu MA, Gannon B. Application of Resource Utilization in Dementia (RUD) instrument in a global setting. *Alzheimers Dement*. 2013;9:429–435.e417.
58. Berg L, Miller JP, Storandt M, Duchek J, Morris JC, Rubin EH, Burke WJ, Coben LA. Mild senile dementia of the Alzheimer type: 2. Longitudinal assessment. *Ann Neurol*. 1988;23:477–484.
59. Hughes CP, Berg L, Danziger WL, Coben LA, Martin RL. A new clinical scale for the staging of dementia. *Br J Psychiatry*. 1982;140:566–572.
60. O'Bryant SE, Lacritz LH, Hall J, Waring SC, Chan W, Khodr ZG, Massman PJ, Hobson V, Cullum CM. Validation of the new interpretive guidelines for the Clinical Dementia Rating Scale Sum of Boxes score in the National Alzheimer's Coordinating Center database. *Arch Neurol*. 2010;67:746–749.
61. Andrews JS, Desai U, Kirson NY, Zichlin ML, Ball DE, Matthews BR. Disease severity and minimal clinically important differences in clinical outcome assessments for Alzheimer's disease clinical trials. *Alzheimers Dement (N Y)*. 2019;5:354–363.
62. FDA. Patient-Focused Drug Development Guidance Public Workshop: methods to identify what is important to patients & select, develop or modify fit-for-purpose clinical outcomes assessments. *Appendix*. <https://www.fda.gov/media/116277/download>. Accessed March 4, 2021
63. Mohs RC, Knopman D, Petersen RC, Ferris SH, Ernesto C, Grundman M, Sano M, Bieliauskas L, Geldmacher D, Clark C, Thal LJ. Development of cognitive instruments for use in clinical trials of antedementia drugs: additions to the Alzheimer's Disease Assessment Scale that broaden its scope. The Alzheimer's Disease Cooperative Study. *Alzheimer Dis Assoc Disord*. 1997;11(Suppl 2):S13–S21.
64. Evans S, McRae-McKee K, Wong MM, Hadjichrysanthou C, De Wolf F, Anderson R. The importance of endpoint selection: how effective does a drug need to be for success in a clinical trial of a possible Alzheimer's disease treatment? *Eur J Epidemiol*. 2018;33:635–644.
65. Kueper JK, Speechley M, Montero-Odasso M. The Alzheimer's Disease Assessment Scale-Cognitive Subscale (ADAS-Cog): modifications and responsiveness in pre-dementia populations. A narrative review. *J Alzheimers Dis*. 2018;63:423–444.
66. Creavin ST, Wisniewski S, Noel-Storr AH, Trevelyan CM, Hampton T, Rayment D, Thom VM, Nash KJ, Elhamoui H, Milligan R, Patel AS, Tsivos DV, Wing T, Phillips E, Kellman SM, Shackleton HL, Singleton GF, Neale BE, Watton ME, Cullum S. Mini-Mental State Examination (MMSE) for the detection of dementia in clinically unevaluated people aged 65 and over in community and primary care populations. *Cochrane Database Syst Rev*. 2016:CD011145.
67. Mitchell AJ. A meta-analysis of the accuracy of the mini-mental state examination in the detection of dementia and mild cognitive impairment. *J Psychiatr Res*. 2009;43:411–431.
68. Murphy KJ, Rich JB, Troyer AK. Verbal fluency patterns in amnesic mild cognitive impairment are characteristic of Alzheimer's type dementia. *J Int Neuropsychol Soc*. 2006;12:570–574.
69. Tabert MH, Manly JJ, Liu X, Pelton GH, Rosenblum S, Jacobs M, Zamora D, Goodkind M, Bell K, Stern Y, Devanand DP. Neuropsychological prediction of conversion to Alzheimer disease in patients with mild cognitive impairment. *Arch Gen Psychiatry*. 2006;63:916–924.
70. Duff K, Clark HJD, O'Bryant SE, Mold JW, Schiffer RB, Sutker PB. Utility of the RBANS in detecting cognitive impairment associated with Alzheimer's disease: sensitivity, specificity, and positive and negative predictive powers. *Arch Clin Neuropsychol*. 2008;23:603–612.
71. Karantzoulis S, Novitski J, Gold M, Randolph C. The Repeatable Battery for the Assessment of Neuropsychological Status (RBANS): utility in detection and characterization of mild cognitive impairment due to Alzheimer's disease. *Arch Clin Neuropsychol*. 2013;28:837–844.
72. Papp KV, Rentz DM, Orlovsky I, Sperling RA, Mormino EC. Optimizing the preclinical Alzheimer's cognitive composite with semantic processing: The PACC5. *Alzheimers Dement (N Y)*. 2017;3:668–677.
73. Wu Y, Zhang X, He Y, Cui J, Ge X, Han H, Luo Y, Liu L, Wang X, Yu H. Predicting Alzheimer's disease based on survival data and longitudinally measured performance on cognitive and functional scales. *Psychiatry Res*. 2020;291:113201.

74. Sikkes SA, Knol DL, Pijnenburg YA, de Lange-de Klerk ES, Uitdehaag BM, Scheltens P. Validation of the Amsterdam IADL Questionnaire®, a new tool to measure instrumental activities of daily living in dementia. *Neuroepidemiology*. 2013;41:35–41.
75. Jutten RJ, Peeters C, Leijdesdorff S, Visser PJ, Maier AB, Terwee CB, Scheltens P, Sikkes S. Detecting functional decline from normal aging to dementia: development and validation of a short version of the Amsterdam IADL Questionnaire. *Alzheimers Dement (Amst)*. 2017;8:26–35.
76. Griffiths AW, Smith SJ, Martin A, Meads D, Kelley R, Surr CA. Exploring self-report and proxy-report quality-of-life measures for people living with dementia in care homes. *Qual Life Res*. 2020;29:463–472.
77. Travis Seidl JN, Massman PJ. Cognitive and functional correlates of NPI-Q scores and symptom clusters in mildly demented Alzheimer patients. *Alzheimer Dis Assoc Disord*. 2016;30:145–151.
78. FDA-NIH Biomarker Working Group. BEST (Biomarkers, EndpointS, and other Tools) Resource. <https://www.ncbi.nlm.nih.gov/books/NBK326791/>. Accessed February, 2021
79. Wu L, Rosa-Neto P, Gauthier S. Use of biomarkers in clinical trials of Alzheimer disease: from concept to application. *Mol Diagn Ther*. 2011;15:313–325.
80. Horgan D, Borisch B, Richer E, Bernini C, Kalra D, Lawler M, Ciliberto G, Van Poppel H, Paradiso A, Riegman P, Triberti S, Metspalu A, Chiti A, Macintyre E, Boccia S, Calvo F, Schatz D, Koeva-Balabanova J, Jonsson B. Propelling health care into the twenties. *Biomed Hub*. 2020;5:15–67.
81. Young PNE, Estarellas M, Coomans E, Srikrishna M, Beaumont H, Maass A, Venkataraman AV, Lissaman R, Jimenez D, Betts MJ, McGlinchey E, Berron D, O'Connor A, Fox NC, Pereira JB, Jagust W, Carter SF, Paterson RW, Scholl M. Imaging biomarkers in neurodegeneration: current and future practices. *Alzheimers Res Ther*. 2020;12:49.
82. Bittner T, Zetterberg H, Teunissen CE, Ostlund RE, Jr., Militello M, Andreasson U, Hubeek I, Gibson D, Chu DC, Eichenlaub U, Heiss P, Kobold U, Leinenbach A, Madin K, Manuilova E, Rabe C, Blennow K. Technical performance of a novel, fully automated electrochemiluminescence immunoassay for the quantitation of β -amyloid (1-42) in human cerebrospinal fluid. *Alzheimers Dement*. 2016;12:517–526.
83. Lifke V, Manuilova E, Knop C, Selle T, Kraus W, Oelschlaegel T, Hillringhaus L. Elecsys® Total-Tau CSF and Elecsys® Phospho-Tau (181P) CSF: novel, fully automated immunoassays for rapid and accurate quantitation of CSF biomarkers for clinical use. Clinical Trials on Alzheimer's Disease (CTAD) 2017, Boston, MA.
84. Hansson O, Seibyl J, Stomrud E, Zetterberg H, Trojanowski JQ, Bittner T, Lifke V, Corradini V, Eichenlaub U, Batrla R, Buck K, Zink K, Rabe C, Blennow K, Shaw LM, Swedish Bio Fsg, Alzheimer's Disease Neuroimaging I. CSF biomarkers of Alzheimer's disease concord with amyloid-beta PET and predict clinical progression: a study of fully automated immunoassays in BioFINDER and ADNI cohorts. *Alzheimers Dement*. 2018;14:1470–1481.
85. F. Hoffman-La Roche Ltd. FDA grants Breakthrough Device Designation for Roche's Elecsys cerebrospinal fluid (CSF) assays to support the improved diagnosis of Alzheimer's disease. <https://www. Roche.com/media/releases/med-cor-2018-07-20.htm>. 20 July, 2018.
86. Sanabria Bóhorquez S, Marik J, Barret O, Tamagnan G, Alagille D, Ayalon G, Ward M, Jennings D, Seibyl J, Marek K, Kerchner GA, Weimer RM. Evaluation of cross-sectional tau burden and preliminary longitudinal changes in Alzheimer's disease subjects using [18F]GTP1 (Genetech tau probe 1). Clinical Trials in Alzheimer's (CTAD) 2015, Barcelona, Spain.
87. ClinicalTrials.gov. U.S. National Library of Medicine. NCT03174938. <https://clinicaltrials.gov/ct2/show/NCT03174938>. Accessed February 2021.
88. Leuzy A, Smith R, Ossenkoppele R, Santillo A, Borroni E, Klein G, Ohlsson T, Jogi J, Palmqvist S, Mattsson-Carlgren N, Strandberg O, Stomrud E, Hansson O. Diagnostic performance of RO948 F18 tau positron emission tomography in the differentiation of Alzheimer disease from other neurodegenerative disorders. *JAMA Neurol*. 2020;77:955–965.
89. Blennow K, Hampel H, Weiner M, Zetterberg H. Cerebrospinal fluid and plasma biomarkers in Alzheimer disease. *Nat Rev Neurol*. 2010;6:131–144.
90. Olsson B, Lautner R, Andreasson U, Öhrfelt A, Portelius E, Bjerke M, Hölltå M, Rosén C, Olsson C, Strobel G, Wu E, Dakin K, Petzold M, Blennow K, Zetterberg H. CSF and blood biomarkers for the diagnosis of Alzheimer's disease: a systematic review and meta-analysis. *Lancet Neurol*. 2016;15:673–684.
91. Blennow K, Zetterberg H. Biomarkers for Alzheimer's disease: current status and prospects for the future. *J Intern Med*. 2018;284:643–663.
92. Blom ES, Giedraitis V, Zetterberg H, Fukumoto H, Blennow K, Hyman BT, Irizarry MC, Wahlund LO, Lannfelt L, Ingelsson M. Rapid progression from mild cognitive impairment to Alzheimer's disease in subjects with elevated levels of tau in cerebrospinal fluid and the APOE epsilon4/epsilon4 genotype. *Dement Geriatr Cogn Disord*. 2009;27:458–464.
93. Hansson O, Zetterberg H, Buchhave P, Londos E, Blennow K, Minthon L. Association between CSF biomarkers and incipient Alzheimer's disease in patients with mild cognitive impairment: a follow-up study. *Lancet Neurol*. 2006;5:228–234.
94. Fagan AM, Roe CM, Xiong C, Mintun MA, Morris JC, Holtzman DM. Cerebrospinal fluid tau/beta-amyloid(42) ratio as a prediction of cognitive decline in nondemented older adults. *Arch Neurol*. 2007;64:343–349.
95. Johnson KA, Fox NC, Sperling RA, Klunk WE. Brain imaging in Alzheimer disease. *Cold Spring Harb Perspect Med*. 2012;2:a006213.
96. Leuzy A, Chiottis K, Lemoine L, Gillberg PG, Almkvist O, Rodriguez-Vieitez E, Nordberg A. Tau PET imaging in neurodegenerative tauopathies-still a challenge. *Mol Psychiatry*. 2019;24:1112–1134.

97. Eli Lilly and Company. Tauvid. Prescribing information. https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/212123s000lbl.pdf. Accessed December 2020.
98. Eli Lilly and Company. Lilly receives U.S. FDA approval of TAUVID™ (flortaucipir F 18 injection) for use in patients being evaluated for Alzheimer's disease. <https://investor.lilly.com/news-releases/news-release-details/lilly-receives-us-fda-approval-tauvidtm-flortaucipir-f-18>. Accessed December 2020.
99. Genentech Inc. Data on file.
100. Wildsmith KR. Tau imaging in anti-tau clinical trials: 12:05 p.m. [18F]GTP1 and the semorinemab biomarker strategy. TAU 2020, Washington, D.C.
101. Sanabria Bóhorquez S, Marik J, Ogasawara A, Tinianow JN, Gill HS, Barret O, Tamagnan G, Alagille D, Ayalon G, Manser P, Bengtsson T, Ward M, Williams SP, Kerchner GA, Seibyl JP, Marek K, Weimer RM. [18F]GTP1 (Genentech Tau Probe 1), a radioligand for detecting neurofibrillary tangle tau pathology in Alzheimer's disease. *Eur J Nucl Med Mol Imaging*. 2019;46:2077–2089.
102. Teng E, Ward M, Manser PT, Sanabria-Bohorquez S, Ray RD, Wildsmith KR, Baker S, Kerchner GA, Weimer RM. Cross-sectional associations between [18F]GTP1 tau PET and cognition in Alzheimer's disease. *Neurobiol Aging*. 2019;81:138–145.
103. Chan D, Janssen JC, Whitwell JL, Watt HC, Jenkins R, Frost C, Rossor MN, Fox NC. Change in rates of cerebral atrophy over time in early-onset Alzheimer's disease: longitudinal MRI study. *Lancet*. 2003;362:1121–1122.
104. Chan D, Fox NC, Scahill RI, Crum WR, Whitwell JL, Leschziner G, Rossor AM, Stevens JM, Cipolotti L, Rossor MN. Patterns of temporal lobe atrophy in semantic dementia and Alzheimer's disease. *Ann Neurol*. 2001;49:433–442.
105. Eskildsen SF, Coupe P, Fonov VS, Pruessner JC, Collins DL, Alzheimer's Disease Neuroimaging Initiative. Structural imaging biomarkers of Alzheimer's disease: predicting disease progression. *Neurobiol Aging*. 2015;36(Suppl 1):S23–S31.
106. McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR, Jr., Kawas CH, Klunk WE, Koroshetz WJ, Manly JJ, Mayeux R, Mohs RC, Morris JC, Rossor MN, Scheltens P, Carrillo MC, Thies B, Weintraub S, Phelps CH. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement*. 2011;7:263–269.
107. Albers MW, Gilmore GC, Kaye J, Murphy C, Wingfield A, Bennett DA, Boxer AL, Buchman AS, Cruickshanks KJ, Devanand DP, Duffy CJ, Gall CM, Gates GA, Granholm AC, Hensch T, Holtzer R, Hyman BT, Lin FR, McKee AC, Morris JC, Petersen RC, Silbert LC, Struble RG, Trojanowski JQ, Verghese J, Wilson DA, Xu S, Zhang LI. At the interface of sensory and motor dysfunctions and Alzheimer's disease. *Alzheimers Dement*. 2015;11:70–98.
108. Kourtis LC, Regele OB, Wright JM, Jones GB. Digital biomarkers for Alzheimer's disease: the mobile/wearable devices opportunity. *NPJ Digit Med*. 2019;2:9.
109. Vernooij-Dassen MJ, Moniz-Cook ED, Woods RT, De Lepeleire J, Leuschner A, Zanetti O, de Rotrou J, Kenny G, Franco M, Peters V, Iliffe S. Factors affecting timely recognition and diagnosis of dementia across Europe: from awareness to stigma. *Int J Geriatr Psychiatry*. 2005;20:377–386.
110. Herrmann LK, Welter E, Leverenz J, Lerner AJ, Udelson N, Kanetsky C, Sajatovic M. A systematic review of dementia-related stigma research: can we move the stigma dial? *Am J Geriatr Psychiatry*. 2018;26:316–331.



Published by
F. Hoffmann-La Roche Ltd
Group Communications
4070 Basel, Switzerland

© 2022

All trademarks mentioned enjoy legal protection.
Elecsys is a trademark of Roche

www.roche.com

M-XX-00008411